

DOE-ER40757-094
 UTEXAS-HEP-97-3

Finite Temperature Wave-Function Renormalization, A Comparative Analysis

Ian A. Chapman

Abstract

We compare two competing theories regarding finite temperature wave-function corrections for the process $H \rightarrow e^+e^-$ and for $n + \nu \rightarrow p + e^-$ and related processes of interest for primordial nucleosynthesis. Although the two methods are distinct (as shown in $H \rightarrow e^+e^-$) they yield the same finite temperature correction for all $n \rightarrow p$ and $p \rightarrow n$ processes. Both methods yield an increase in the He/H ratio of .01% due to finite temperature renormalization rather than a decrease of .16% as previously predicted.

1 Introduction

To calculate finite temperature corrections to physical processes beyond the tree level, one needs to have a consistent and correct method of temperature dependent renormalization. Unfortunately, there seems to be more than one way to do this. Although everyone seems to agree on how to handle temperature dependent phase space, temperature dependent vertex corrections, effective mass corrections, and temperature dependent bremsstrahlung corrections, they do not agree on how one handles wave function renormalization.

In this paper, we will examine two competing models. Each model's effect on the 'effective' projection operator (which in turn will tell us how to handle wave function renormalization) is examined for the general case. These

general results are then applied to $H \rightarrow e^+e^-$ since this is a fairly simple process to understand. We also apply both methods to processes of interest for primordial nucleosynthesis, specifically $n + \nu \rightarrow p + e^-$ and related processes. These processes were first analyzed by Dicus, et al.[1], and independently by Cambier, Primack, and Sher[2]. Since these papers were published, however, it has been claimed that neither paper handled the finite temperature wave function renormalization scheme correctly. Both Donoghue and Holstein[3], and Sawyer[4] propose models that claim to correct this difficulty. Both agree that one must use *finite temperature* Dirac spinors (and thus projection operators), but otherwise their models seem to differ significantly.

Before we begin a few comments about gauges are in order. The correction of interest will be the wavefunction renormalization part of the width. Since this is a gauge dependent quantity, we might wonder if our choice of gauge will make a difference in our results. We solve this problem in two ways. For most of this paper, we will avoid a specific choice of gauge. Instead we will discuss the differences between the models in the most general way possible. When we calculate a specific result for nucleosynthesis, this is no longer a viable option. In Appendix I, however, we prove that we can talk about the differences in the wavefunction renormalization parts in the width in a gauge invariant manner. We take advantage of this in Eq.(14).

2 Generalized Analysis

As discussed in the previous literature, the presence of a background thermal bath breaks Lorentz invariance. At $T=0$ we can identify the wavefunction renormalization constant Z_2 for fermions with momentum p as $1-Z_2 = \left. \frac{\partial \Sigma}{\partial \not{p}} \right|_{\not{p}=m}$ where Σ is the self energy. Since Lorentz invariance is broken when $T \neq 0$, it is claimed that this identification of a momentum independent renormalization constant is no longer correct. Donoghue and Holstein advise us to look at the modified propagator instead and to modify the Dirac spinors accordingly. Thus the modified propagator becomes

$$G = \frac{1}{\not{p} - m - \Sigma} \quad (1)$$

where the fermion self-energy, Σ is

$$\Sigma = A p_0 \gamma^0 - B p_i \gamma^i - C. \quad (2)$$

A, B , and C are momentum dependent scalar functions. We identify an "effective" momentum, $\tilde{p}^\mu$ where $\tilde{p}^\mu = ((1 - A(p))p^0, (1 - B(p))p^i)$ and an effective mass $\tilde{m}$ where $\tilde{m} = m - C(p)$. We are instructed to replace the fermion momenta and mass with the "effective" fermion momenta and mass everywhere inside the Dirac spinors. When this is done, *and* the projection operator is expanded about the *physical* mass shell energy, we get the following "effective" projection operator

$$\Lambda_{eff}^+ = \frac{\not{\tilde{p}} + \tilde{m}}{2\tilde{p}_0} \quad (3)$$

Converting this into the mass shell p^μ , expanding about the mass-shell energy, and writing this in terms of A, B , and C , we get (noting that $\omega_p = \sqrt{\mathbf{p}^2 + m^2}$ or the mass shell energy)

$$\Lambda_{eff}^+ = \left(1 - \frac{\partial((1-2A)p_0^2 - (1-2B)\mathbf{p}^2 - (1-\frac{2C}{m})m^2)}{\partial p_0} \bigg|_{p_0=\omega_p} \right) \left(\frac{\not{p} + m}{2\omega_p} \right) \quad (4)$$

Simplifying this, we find

$$\Lambda_{eff}^+ = \left(A + \frac{\partial A}{\partial p_0} \bigg|_{p_0=\omega_p} \omega_p - \frac{\partial B}{\partial p_0} \bigg|_{p_0=\omega_p} \frac{\mathbf{p}^2}{\omega_p} - \frac{\partial C}{\partial p_0} \bigg|_{p_0=\omega_p} \frac{m}{\omega_p} \right) \left(\frac{\not{p} + m}{2\omega_p} \right) \quad (5)$$

Thus the effective projection operator is equal to the original $T=0$ projection operator multiplied by a single scalar momentum dependent function. Donoghue and Holstein identify this function as $1 - Z_2$ where " Z_2 " is the effective temperature dependent wave-renormalization factor.

Sawyer gives us a differing way to handle finite temperature wave-function renormalization. Sawyer tells us to start with the same modified propagator as shown in Eq.(1), but then reminds us that the propagator can also be written in terms of the effective Dirac spinors as shown by Weldon[5]. Thus

$$G = \frac{\sum_\alpha w_\alpha \bar{w}^\alpha}{p_0 - \omega_p} + \text{finite terms} \quad (6)$$

where w is the temperature dependent renormalized Dirac spinor. Using this fact, Sawyer is able to perturbatively derive the temperature dependent

renormalized Dirac spinors, w , in terms of the unrenormalized zero temperature Dirac spinors, u . He was then able to use these new spinors to directly define the effective projection operators. If we follow this procedure we get

$$\Lambda_{eff}^+ = \frac{1}{4\omega_p} (\gamma^0 \frac{\partial \Sigma}{\partial p_0} \Big|_{p_0=\omega_p} (\not{p} + m) + (\not{p} + m) \frac{\partial \Sigma}{\partial p_0} \Big|_{p_0=\omega_p} \gamma^0) \quad (7)$$

When we write Sawyer's effective projection operator in terms of A, B, C , and ω_p as defined previously, we get (evaluating p_0 at ω_p).

$$\begin{aligned} \Lambda_{eff}^+ = & \left(A + \frac{\partial A}{\partial p_0} p_0 \right) \left(\frac{\not{p} + m}{2\omega_p} \right) \\ & + \left(\frac{\partial B}{\partial p_0} p_i \gamma^i p_0 - \frac{\partial B}{\partial p_0} \mathbf{p}^2 \gamma^0 - \frac{\partial C}{\partial p_0} p_0 - \frac{\partial C}{\partial p_0} m \gamma^0 \right) \left(\frac{1}{2\omega_p} \right) \end{aligned} \quad (8)$$

It is immediately evident that the above expression for Λ_{eff}^+ is in general different from the expression for Λ_{eff}^+ in Eq.(5). Specifically, we note that Λ_{eff}^+ is *not* equal to the unrenormalized projection operator times some scalar functional factor. We see this explicitly in the process $H \rightarrow e^+ e^-$.

3 $H \rightarrow e^+ e^-$ comparison

The tree-level matrix element for this process in momentum space is

$$M_T = -\frac{igm}{2M_W} \bar{u}(e^-) v(e^+) \quad (9)$$

where m is the mass of the fermion, M_W is the mass of the $W^\pm$, $\bar{u}$ and v are the Dirac Spinors at $T=0$, and M_T is the tree level matrix element for this process. The modified width considering *only* those terms due to the temperature dependent wavefunction renormalization can be found (to first order in α) by cross-multiplying the tree-level matrix elements using the modified projection operators, Λ_{eff}^+ and Λ_{eff}^- , in place of the normal zero temperature unrenormalized projection operators. One then integrates over the correct temperature dependent phase space. Since this later part is not in dispute, it is easier to simply compare the modified spin-summed squared self-energy matrix elements in both models. (As I mentioned before, I will

neglect the temperature dependent mass terms here.) We assume for this process that the Higgs is at rest with respect to the background thermal bath, and that the Higgs is significantly more massive than the background temperature. When we use the modified projection operator for Donoghue and Holstein's model, we get

$$|M|^2 = \frac{2M_H^2 g^2 m^2 (1 - \frac{4m^2}{M_H^2})}{4M_W^2} \left(A + \frac{\partial A}{\partial e_0} e_0 - \frac{\partial B}{\partial e_0} \frac{\mathbf{e}^2}{e_0} - \frac{\partial C}{\partial e_0} \frac{2m}{M_H} \right) \quad (10)$$

When we use the modified projection operator as described by Sawyer, however, we get

$$|M|^2 = \frac{g^2 m^2}{4M_W^2} \left[2M_H^2 \left(1 - \frac{4m^2}{M_H^2} \right) \left(A + \frac{\partial A}{\partial e_0} e_0 \right) + \frac{\partial C}{\partial e_0} m M_H \right] \quad (11)$$

These matrix elements squared yield differing widths when integrated over phase space. Thus the two models are distinct for $H \rightarrow e^+ e^-$.

4 $n + \nu \rightarrow p + e^-$ comparison

Finite temperature corrections have been of considerable interest for $n \rightarrow p$ and $p \rightarrow n$ reactions because of nucleosynthesis. Specifically, one would like to know how finite temperature corrections affect abundance ratios such as H/He in the early universe. Since all $n \rightarrow p$ and $p \rightarrow n$ processes are related by virtually identical matrix elements, a comparison of $n + \nu \rightarrow p + e^-$ is sufficient for all such processes.

In this comparison we make the following standard assumptions (valid for proton to neutron ratios in the early universe). First we assume that the ambient temperatures of interest are of order 1 MeV so that the proton and neutron have considerably more mass than the ambient temperature. (Thus any terms of the form $\frac{m}{M_p}$ or $\frac{m}{M_n}$ are ignored.) We also assume that the initial nucleon is at rest with respect to the surrounding medium. The neutrino electron part of the spin-summed squared matrix element looks like

$$\text{Tr} \left\{ \Lambda_{eff}^+ \frac{1}{2} \gamma^\mu (1 - \gamma^5) \not{p} \frac{1}{2} (1 + \gamma^5) \gamma^\nu \right\} \quad (12)$$

In general Λ_{eff}^+ has three kinds of terms. There are scalar terms, terms that are multiplied by a factor of γ^0 , and the remaining terms which are all

multiplied by a factor of $e_i \gamma^i$. The scalar terms will trace away, and those terms that are multiplied by $e_i \gamma^i$ do not survive the phase space integration. Thus the *only* terms in the effective projection operators that matter are those terms which are multiplied by γ^0 . We now look back at Donoghue and Holstein's effective projection operator (Eq.(5)) and Sawyer's effective projection operator (Eq.(8)). Isolating the γ^0 terms in both models, we get

$$\Lambda_{eff}^+ = \left(A + \frac{\partial A}{\partial e_0} e_0 - \frac{\partial B}{\partial e_0} \frac{e_v^2}{e_0} - \frac{\partial C}{\partial e_0} \frac{m}{e_0} \right) e_0 \gamma^0 \quad (13)$$

i.e., the effective projection operators are *identical*. In the above equation, e_v is the magnitude of the 3-momentum of the electron. To verify this, I did this calculation using Donoghue and Holstein's method *and* Sawyer's method using the Feynman gauge. (Sawyer used the Coulomb gauge in his analysis.)

When we calculate the width according to Sawyer (and Donahue and Holstein) minus the width according to Dicus et al. (and Cambier et al.) for the reaction $n + \nu \rightarrow p + e^-$ we get

$$\begin{aligned} \delta\Gamma_{Sawyer} - \delta\Gamma_{Dicus} = & \frac{G_F^2 (g_V^2 + 3g_A^2) \alpha}{2\pi^4} \int_Q (Q - e)^2 e_v de \int_0 k^2 dk \\ & [1 + \exp(-e/T)]^{-1} \left[1 + \exp\left(\frac{e - Q}{T}\right) \right]^{-1} \times \left\{ \frac{n_f(E_k)}{E_k} \left[\frac{4m^2(e + E_k)}{(E_k e + m^2)^2 - e_v^2 k^2} \right. \right. \\ & + \frac{4m^2(e - E_k)}{(E_k e - m^2)^2 - e_v^2 k^2} + \frac{E_k}{e_v k} \text{Ln} \left| \frac{(E_k e - e_v k)^2 - m^4}{(E_k e + e_v k)^2 - m^4} \right| \\ & \left. \left. + \frac{e}{e_v k} \text{Ln} \left| \frac{(E_k^2 e^2 - (e_v k - m^2)^2)}{E_k^2 e^2 - (e_v k + m^2)^2} \right| \right] - 2 \frac{n_b(k) \text{Ln} \left| \frac{e + e_v}{e - e_v} \right|}{e_v k} \right\} \quad (14) \end{aligned}$$

where Q is the rest mass of the neutron minus the rest mass of the proton, $e_v = \sqrt{e^2 - m^2}$, $E_k = \sqrt{k^2 + m^2}$, $n_f(E_k) = [1 + \exp(E_k/T)]^{-1}$, $n_b(k) = [\exp(k/T) - 1]^{-1}$. Using the above equation and its analogs for the other five $n \rightarrow p$ and $p \rightarrow n$ reactions, we obtain the change in the width ratios in figures one and two.

In figure three we combine the $n \rightarrow p$ and $p \rightarrow n$ processes into a single ratio change for $n \rightarrow p$ and $p \rightarrow n$.

We use these results to solve for the change in the asymptotic neutron to baryon ratio, δX_n as described by Bernstein[6] where

$$\delta X_n(t) = S(t) \int^t dt' [\delta\Gamma(t')_{p \rightarrow n} [1 - X_n(t')] - \delta\Gamma(t')_{n \rightarrow p} X_n(t')] S^{-1}(t') \quad (15)$$

where $S(t)$ is the solution of $\frac{dS}{dt} + (\Gamma_{n \rightarrow p} + \Gamma_{p \rightarrow n})S = 0$.

We solve Eq.(15) for the δX_n change we expect. We solve for $S(t)$, $\delta\Gamma_{p \rightarrow n}$, and $\delta\Gamma_{n \rightarrow p}$ using Vegas (a Monte-Carlo numeric integration routine). We solve the outer integration over t by Gaussian quadratures. We obtain an increase of δX_n of 0.000415. Thus in Dicus et al.'s original paper, $Y_3 - Y_4$ in Table 1 (on page 2701) should be changed from +0.0004 to -0.000015. This corresponds with an increase of .01% in δX_n due to finite temperature renormalization as compared to a .16% decrease as predicted by Dicus et al. and Cambier et al. Sawyer predicts an increase of .02% in δX_n due to finite temperature renormalization. Although his result differs somewhat from ours, we both agree that the change in δX_n should be very small and positive i.e., increase due to temperature dependent renormalization effects.

5 Summary

It is clear that the models proposed by Donoghue and Holstein, and Sawyer are indeed *distinct*. However, it is equally clear that both models yield the same finite temperature corrections for $n \rightarrow p$ and $p \rightarrow n$ processes when the temperature is much less than the nucleon mass. This is because the later processes, at least as described in the early universe, are insensitive to any correction that does not depend directly on the γ^0 or energy component of the effective projection operator. It is extremely interesting to note that the effective projection operators for both models are *identical* in this one crucial component.

If one can show that in one's process of interest, the γ^i and scalar pieces of the renormalized projection operator don't matter, then you can indeed use Donoghue and Holstein's effective "Z₂" in the conventional way. Otherwise, it is not clear which method, if either, should be used.

6 Acknowledgments

I especially wish to thank D.A. Dicus for his assistance. Without his patience, insight, support, and encouragement, this research would not have been possible. I also thank M. Byrd and X. Bonnin for discussions and technical assistance. This research was supported in part by the U.S. Department of Energy under Contract # DE-FG3-93ER40757.

7 Appendix I

We want a gauge independent way of comparing two finite temperature renormalization models. It is not clear that looking solely at the wavefunction or self-energy part of the renormalized width will be sufficient since this part is gauge dependent. Fortunately we know that all of the overall models we consider in this paper are gauge invariant *and* that the only part of these models that differ are the self-energy parts. Thus

$$\delta\Gamma_{Sawyer}^{total} = \delta\Gamma_{Sawyer}^{SE} + \delta\Gamma_{Sawyer}^{other} \quad (16)$$

and

$$\delta\Gamma_{Dicus}^{total} = \delta\Gamma_{Dicus}^{SE} + \delta\Gamma_{Dicus}^{other} \quad (17)$$

In the above equations, $\delta\Gamma^{total}$ represents the total change in width due to finite temperature renormalization, $\delta\Gamma^{SE}$ represents the change in width due to finite temperature wavefunction renormalization, while $\delta\Gamma^{other}$ represents all other changes in the width due to finite temperature renormalization. Since the models proposed by Dicus and Sawyer *only* disagree on the wavefunction renormalization part, we conclude that $\delta\Gamma^{other}$ must be the same for both models. Thus if we know subtract Eq.(17) from Eq.(16) we get

$$\delta\Gamma_{Sawyer}^{total} - \delta\Gamma_{Dicus}^{total} = \delta\Gamma_{Sawyer}^{SE} - \delta\Gamma_{Dicus}^{SE} \quad (18)$$

Since we *know* the total change in widths is gauge independent for both models, it follows from Eq.(18) that $\delta\Gamma_{Sawyer}^{SE} - \delta\Gamma_{Dicus}^{SE}$ must be gauge independent as well. This gives us a good way to compare two different models for finite temperature renormalization.

References

- [1] D.A. Dicus, E.W. Kolb, A.M. Gleeson, E.C.G. Sudarshan, V.L. Teplitz, M.S. Turner, *Physical Review* **D26**, 2694-2706 (1982)
- [2] J-L. Cambier, J.R. Primack, and M. Sher, *Nuclear Physics* **B209**, 372-388 (1982)
- [3] J.F. Donoghue and B.R. Holstein, *Physical Review* **D28**, 340-348 (1983); *Physical Review* **D29**, 3004 (1984)
- [4] R.F. Sawyer, *Physical Review* **D53**, 4232-4236 (1996)
- [5] H.A. Weldon, *Physical Review* **D40**, 2410-2420 (1989)
- [6] J. Bernstein, L.S. Brown, G. Feinberg, *Review of Modern Physics* **61**, 25 (1989)




Figure 1: These are the ratios of the change of the finite temperature first order corrections (as calculated in Eq.(14)) to the finite temperature uncorrected width for all $n \rightarrow p$ processes in units of $\frac{\alpha}{\pi}$. The temperature is measured in units of the electron mass. The short dashed line represents $n \rightarrow p e \nu$, the long dashed line represents $n e \rightarrow p \nu$, and the solid line represents $n \nu \rightarrow p e$.




Figure 2: These are the ratios of the change of the finite temperature first order corrections (as calculated in Eq.(14)) to the finite temperature uncorrected widths for all $p \rightarrow n$ processes in units of $\frac{\alpha}{\pi}$. The temperature is measured in units of the electron mass. The short dashed line represents $p\nu \rightarrow ne$, and the long dashed line represents $pe \rightarrow n\nu$




Figure 3: These are the combined changes in the finite temperature first order corrections (as calculated in Eq.(14)) to the combined finite temperature uncorrected widths for both $n \rightarrow p$ and $p \rightarrow n$ in units of $\frac{\alpha}{\pi}$. The temperature is measured in units of the electron mass. The solid line represents the combined $n \rightarrow p$ processes while the dashed line represents the combined $p \rightarrow n$ processes.